

Antihyperglycemic and Antihyperlipidemic Effects of Methanolic Seeds Extract of Pandanus odoratissimus in Alloxan-Induced Diabetic Rats

by Ernawati Sinaga

Submission date: 09-Feb-2021 11:29AM (UTC+0700)

Submission ID: 1505154474

File name: B9-Endarti-erna-2020.pdf (874.14K)

Word count: 6096

Character count: 33093

1 Antihyperglycemic and Antihyperlipidemic Effects of Methanolic Seeds Extract of *Pandanus odoratissimus* in Alloxan-Induced Diabetic Rats

Sri Endarti Rahayu¹, 52 awati Sinaga¹, Suprihatin¹, Arief Dwi Jayanto¹, Yedi Adistron¹

¹Faculty of Biology, Universitas Nasional, Jl. Sawo-Manila, South Jakarta 12520, Indonesia

*Corresponding author: endarti@civitas.unas.ac.id

2 ABSTRACT

Diabetes mellitus is one of the most serious diseases affecting a larger global population, accompanied with other long-term complications. Also, the management of this disease comes with various side effects, hence, the need for substances with little or no side effect. Therefore, this study aimed at evaluating the antidiabetic and antihyperlipidemic effects of methanolic extract of *Pandanus odoratissimus* seeds in alloxan-induced diabetic rats. 30 adult male rats (Sprague Dawley) weighing 180-220 g randomly selected and placed into 6 groups of 5 animals each. The alloxan-induced diabetic rats were orally supplemented with three doses of *P. odoratissimus* methanolic seed extracts, 300, 600 and 900 mg/kg bw, or 0.5 mg/kg of glibenclamide for 2 weeks. The glucose, HbA1c, and lipid profile were measured from the blood sample taken from all studied groups. The results were expressed as mean ± SEMs and compared with repeated measurement using SPSS Version 19.0. The methanolic seed extract of *P. odoratissimus* significantly reduced ($p < 0.05$) the blood glucose level, with a high dose producing 79.15% reduction after 14 days, compared with the control and glibenclamide-treated groups. Furthermore, the seeds extract significantly reduced the concentration of HbA1c, TC, TG and LDL while increased HDL in alloxan-induced diabetic rats. The *P. odoratissimus* could be used to effectively manage diabetes and other related complications.

Keywords: *Pandanus odoratissimus*, seed, methanolic extract, alloxan, glibenclamide.

Correspondance:
Sri Endarti Rahayu¹

¹Faculty of Biology, Universitas Nasional, Jl. Sawo-Manila, South Jakarta 12520, Indonesia

*Corresponding author: endarti@civitas.unas.ac.id

INTRODUCTION

Diabetes is a long-term disease affecting the lives and well-being of individuals, families, and societies at large. This chronic condition is among the top 10 causes of death in adults, and estimated to have caused over four million deaths globally in 2017 (1). Diabetes mellitus is a complex disorder characterized by hyperglycemia due to malfunction in insulin secretion, caused by impaired metabolism of glucose, lipids, and protein (2). Its complication is linked with oxidative stress induced by hyperglycemia suppressing the body's natural antioxidant system (3). In the later stages of this condition, lipid metabolism is affected, in the form of hyperlipidemia and hypercholesterolemia, which are risk factors in atherosclerosis (4). Also, the frequency of hyperlipidemia in diabetic patients is usually very high, depending on the type of diabetes and its degree of control (5).

The disease is managed through insulin injection and oral administration of hypoglycemic drugs such as sulfonylureas and biguanides (6). However, most antidiabetic medications usually come with various side effects, such as liver problem, lactic acidosis, diarrhea and other secondary issues (7). Managing the disease without any side effects is still a challenge in the medical system. This has led to an increase in demand for natural products with anti-hyperglycemic property having little or no side effects. Hence, natural drugs are receiving more attention due to their safety. Thus, plants with antidiabetic activities are vital in the development of diabetes drugs (8). Also, there are many medicinal herbs which have been recommended for the treatment of diabetes (9).

One of these medicinal plants is *Pandanus odoratissimus* which is member of Pandanaceae family. It is a coarsely branched tree, with open crown and aerial roots close to its base. This plant is popularly known as "screw pine" and locally called "pandan samak". It is widely distributed in Indonesia and commonly found on sandy beaches, littoral thickets, on the edges of brackish marshes and mangrove, and inland along watercourses with low attitudes (10) (11). The leaves of *P. odoratissimus* have been reported to be useful for preventing epilepsy (12); infertility (13); inflammation (14) and as CNS-depressant (15). Also, its roots have been reported to have anticancer (16); and antidiabetic effects in patients (17); the flowers are antidiabetic (18), while its fruits are useful as anti-inflammatory and analgesic substances (19). The active component of the leaves extract include alkaloid, flavonoid and saponin (13) while alkaloid, glycosides, steroid, tannin and phenol are the active component of the fruit extract (19).

According to Tiwari and Rao (20) and Hakim et al (21), plants with flavonoid, alkaloid and glycosides possess some antioxidant and antidiabetic properties. Flavonoid reproduce the damaged cell of pancreas and saponin while polyphenol compounds decrease the transportation of glucose transport by suppressing sodium glucose co-transporter 1 (SGLUT) in the intestine. Similarly, Yang et al (22) reported that flavonoid and polyphenol have lipid lowering effects by stimulating glucose utilization in peripheral tissues. There are limited research on the seed of *P. odoratissimus*, as previous studies on food on the leaves, roots, flowers and fruits. Therefore, this study aimed at evaluating the antidiabetic and antihyperlipidemic effect of methanolic extract of *P. odoratissimus* seeds and the

biochemical parameters for the management of diabetes in alloxan-induced rats.

MATERIALS AND METHODS

Plant Material

The *P. odoratissimus* seeds were collected from Calang village, Aceh Jaya district, Aceh Province. The plant fruits from which the seeds were collected were mature, shown by the orange color of its exoderm. These seeds were shade dried, and mechanically grinded to a powder form.

Preparation of Methanolic Extract of *Pandanus odoratissimus* Seed

About 1000 g of the seed powder were subjected to cold maceration using 98% methanol for 15 days, while changing the solvent every 5 days. The collected solvent were filtered with whatman paper and then evaporated at 40°C under low pressure till a semisolid residue was obtained.

Preliminary Phytochemical Screening

The qualitative phytochemical evaluation of the methanolic extracts was conducted to determine the presence or absence of flavonoids through Shinoda test, sterols through Libermann Buchard test, phenols through ferric chloride test, alkaloids through Dragendorff test, and saponins through saponification test, as described by (26).

Antioxidant Activity

DPPH Radical Scavenging Activity

The free radical scavenging capacity of the extracts was determined using DPPH (27). DPPH solution (0.004% w/v) was prepared in 99% methanol, which was then mixed with the seeds extracts to prepare the stock solution of 5 mg/ml. Freshly prepared DPPH solution (0.004% w/v) was dispensed into test tubes after which the seed extract was added. This was followed by the serial dilution (1 to 500 µg) in each test tube to a final volume of 3 ml, then the absorbance read at 515 nm using a spectrophotometer after 10 minutes. Ascorbic acid was used as a reference standard and dissolved in distilled water to make a stock solution of 1 mg/ml. The control sample prepared has the same volume but with no extract and ascorbic acid, as the 95% methanol was served as blank. Then, the percentage scavenging of the DPPH free radical was measured using the equation:

$$\% \text{ Scavenging Activity} = \frac{\text{Absorbance of the control} - \text{Absorbance of the test sample}}{\text{Absorbance of the control}} \times 100$$

The inhibition curve was plotted for duplicate experiments and represented as percentage of mean inhibition \pm standard deviation. IC₅₀ values were obtained by probit analysis (28).

Experimental Animals

Adult male sprague Dawley weighing 180-220 g were used in this study. The animals were kept in cages with standard laboratory condition (temperature 22-23°C with a 12/12 hr light-dark cycle), in animal house of Faculty of Biology, Universitas Nasional Jakarta, Indonesia. Food and water were the available ad libitum. The experimental procedure was approved by the Ethic Committee of Health research from Universitas Pembangunan Nasional "Veteran Jakarta, Indonesia No. B/1719/3/2019/KEPK.

68

Induction of Diabetes Mellitus

The animals were allowed to acclimatize for one week and diabetic models were made by chemically inducing the rats with freshly prepared solution of alloxan monohydrate dissolved in NaCl 0.9%, at a dose of 125 mg/kg body weight. This was injected by intraperitoneally, then the blood glucose level of the rats with hyperglycemia were measured through the tail vein using glucometer. The concentration of glucose level above 200 mg/dl was considered in the experiment.

13

Experimental Design

All animals were randomly divided into six group (n=5) and treated daily for 2 weeks as follows:

Group I (CN) : made up of normal rats, orally given water and food.

Group II (DM): made up of diabetic rats which received alloxan 125 mg/kg b.w through intraperitoneal injection

Group III (DM+Gliben):made up of diabetic rats were received glibenclamide 5 mg/kg b.w through intraperitoneal injection

Group IV (DM+ MEPOS 300) : made up of diabetic rats orally given *P. odoratissimus* methanolic seed extract 300 mg/kg bw once daily for 14 days

Group V (DM+ MEPOS 600) : made up of diabetic rats orally given *P. odoratissimus* methanolic seed extract 600 mg/kg bw once daily for 14 days

Group VI (DM+ MEPOS 900) : made up of diabetic rats orally given *P. odoratissimus* methanolic seed extract 900 mg/kg bw once daily for 14 days

49

Determination of Blood Glucose Levels

Blood samples were collected through the rats' retro orbital plexus to know the blood glucose levels at intervals of 0, 7 and 14 days. The actual glucose level was measured through Achu-check glucometer and reported as mg/dl.

Serum Biochemical Analysis

Glycosylated hemoglobin (HbA1c) levels were determined through the anion exchange HPLC methods using Biored D10, Thiruvananthapuram, India, in accordance with standard protocols described by the manufacturer. The serum total cholesterol (TC) levels were estimated using colorimetric enzymatic methods using CHOD-PAP, Dialab GmbH, Viena, Austria, using the standard protocols described by manufacturer. Serum triglycerides (TG), low-density lipoprotein-cholesterol (LDL-C) and high density lipoprotein-cholesterol (HDL-C) levels were all estimated using random access chemistry analyzer BT 3500, Rome, Italy, based on the standard protocols described by manufacturer.

22

Statistical Analysis

All the relevant data were analyzed with One-Way ANOVA from SPSS 19.0, USA, followed by Tukey's multiple comparison test. Then, *p* values less than 0.05 (*p*< 0.05) were considered significant.

3 RESULT

Phytochemical Screening

Phytochemical screening of the methanolic extract of *P. odoratissimus* seed tested positive for flavonoids, tannins,

saponins, and quinons, but tested negative for alkaloid, steroid and triterpenoid, as shown in Table 1.

Table 1. Phytochemical screening of *Pandanus odoratissimus* seed extract

Phytochemicals	<i>P. odoratissimus</i>
Flavonoids	+
Alkaloids	-
Tannins	+
Saponins	+
Quinons	+
Steroid	-
Triterpenoid	-

Presence of phytochemicals is denoted by (+) sign, while the absence is denoted by (-) sign.

16 Antioxidant Activity

The antioxidant activity of the seed extract was evaluated using *in vitro* assays namely DPPH. The concentration up to 50% inhibition (IC50) is given in Table 2. The antioxidant activities were compared with ascorbic acid which were found

in (29) and the better activity was reflected by lower IC50 values. The *P. odoratissimus* seed extract exhibited significant antioxidant activity, IC= 31.25 µg/ml, which was found to be more potent than those of reference antioxidant, ascorbic acid, with IC50 = 73.61 µg/ml.

Table 2. Antioxidant properties of crude extract of *Pandanus odoratissimus* seed

Sample	Radical scavenging activity IC50 (µg/ml)
<i>Pandanus odoratissimus</i>	31.25
Ascorbic Acid	73.61 [29]

74 Antidiabetic Effect of Methanolic Extract from *Pandanus odoratissimus* Seed

The effect of *P. odoratissimus* seed methanolic extract on blood glucose level in alloxan-treated rats is presented in Table 3. The injection of alloxan induced a significant increase ($p < 0.05$) in blood glucose level in this group at 418.50 ± 113.65 mg/dl, compared with normal control group 97.75 ± 5.44 mg/dl. However, the methanolic extract showed a continuous significant reduction in the glucose level ($p < 0.05$), particularly 14 days after treatment in the diabetic rats. Also, low dosage of the extract significantly reduced ($p < 0.05$) the glucose level from 496.000 ± 76.95 mg/dl to 338.50 ± 67.36 mg/dl at day 7 and to 145.75 ± 36.67 mg/dl at day 14, in

comparison with untreated rat at 418.50 ± 113.65 mg/dl. In addition, this effect obtained with the lower dose is consistent with glibenclamide, which was initially at 946.250 ± 112.82 mg/dl, reduced to 241.25 ± 29.74 mg/dl at day 7, and then to 130.00 ± 14.72 mg/dl at day 14. Similarly, treatment with a high dose of the seed extract (900mg/kg bw) caused a maximum reduction in blood glucose level, from 507.75 ± 72.62 mg/dl to 131.50 ± 13.50 mg/dl at day 7 and then to 104.75 ± 4.50 mg/dl at day 14, compared with the normal control of 97.75 ± 5.66 mg/dl at day 7 and 101.50 ± 8.35 mg/dl at day 14. Furthermore, this effect was more potent than those of the reference drug.

Table 3. Effect of MEPOS on blood glucose and HbA1c in alloxan-induced diabetic rat

Group	Fasting blood glucose (mg/dL)			HbA1c (%)
	Day 0	Day 7	Day 14	
Normal control	97.75 ± 5.44 [#]	96.00 ± 5.66 [#]	101.50 ± 8.35 [#]	6.50 ± 0.12 [#]
Diabetic control	418.50 ± 113.65	462.50 ± 97.43	499.75 ± 146.04	11.75 ± 0.87
Glibenclamide	462.50 ± 112.82	241.25 ± 29.74*	130.00 ± 14.72*	6.50 ± 0.34*
MEPOS (300 mg/kg)	496.00 ± 76.95	338.50 ± 67.36*	145.75 ± 36.67*	9.38 ± 0.12*
MEPOS (600 mg/kg)	505.50 ± 109.00	222.50 ± 35.09*	110.75 ± 9.50*	7.68 ± 0.48*
MEPOS (900 mg/kg)	507.75 ± 72.62	131.50 ± 13.50*	104.75 ± 4.50*	6.32 ± 0.15*

All values are Mean ± SE. # $p < 0.05$ as compare with normal control * $p < 0.05$ as compare to diabetic control. SE : Stabdard Error ; MEPOS : Methanolic extract of *Pandanus odoratissimus* seed

56
Table 4. Effect of MEPOS on lipid profile in alloxan-induced diabetic rat

Groups	Serum lipid profile (mg/dL)			
	TCH	TG	LDL	HDL
Normal control	68.25 ± 5.67 [#]	74.25 ± 6.23 [#]	10.25 ± 0.95 [#]	45.25 ± 1.50 [#]
Diabetic control	88.75 ± 5.25	191.25 ± 3.86	22.25 ± 2.98	32.25 ± 2.06
Glibenclamide	75.50 ± 2.64	85.50 ± 8.81	18.25 ± 1.89	36.50 ± 2.38
MEPOS (300 mg/kg)	67.50 ± 1.29*	110.75 ± 2.50*	13.75 ± 0.50*	41.50 ± 1.73*
MEPOS (600 mg/kg)	61.50 ± 1.73*	93.50 ± 3.10*	11.25 ± 0.95*	46.25 ± 1.25*
MEPOS (900 mg/kg)	52.75 ± 1.70*	74.75 ± 2.75*	10.50 ± 0.57*	56.75 ± 5.31*


27 All values are Mean ± SE. #p<0.05 as compare with normal control *<0.05as compare to diabetic control. SE : Stabdard Error : MEPOS : Methanolic extract of *Pandanus odoratissimus* seed

60
Effect of *Pandanus odoratissimus* Seed Extract on Glycosylated Hemoglobin in Alloxan-Induced Diabetic Rats

The data in Table 3 show that the HbA1c% levels of untreated diabetic rats highly increased to 180.8% of normal control, while treatment with *P. odoratissimus* seed extract 300, 600, 900 mg/kg or glibenclamide reduced this elevation, where the HbA1c% levels were 144.3; 118.1; 97% and 100% respectively.

67
Effect of *Pandanus odoratissimus* Seed Extract on Lipid Profile in Alloxan-Induced Diabetic Rats

Table 4 shows the summary of the serum lipid profile, serum total cholesterol (TC), total triglycerides (TG), low density lipoprotein (LDL) and high density lipoprotein (HDL). There was a significant increase ($p<0.05$) in the level of serum TC, TG and LDL and significant decrease ($p<0.05$) in the level serum HDL of diabetic untreated rats, compared with the control. However, diabetic rats treated with *P. odoratissimus* seed extract had a reversal in the serum lipid profile to almost normal levels with the highest dose (900 mg/kg body weight) being the most effective.



84
8
Figure 1. Effect of different dose of *Pandanus odoratissimus*, libenclamide and their combination on fasting plasma glucose on alloxan induce diabetes rat two weeks treatment. Values are represented as mean ± SEM; n=5, #p<0.05 as compare with normal control *<0.05as compare to diabetic control. Here, NC (Normal Control), DC (diabetic control), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 300 mg/kg bw), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 600 mg/kg bw), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 900 mg/kg bw).




Figure 2. Effect of different dose of *Pandanus odoratissimus*, Glibenclamid and their combination on fasting plasma glucose on alloxan induce diabetes rat two weeks treatment. Values are represented as mean \pm SEM; n=5, #p<0.05 as compare with normal control *<0.05as compare to diabetic control. Here, NC (Normal Control). DC (diabetic control), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 300 mg/kg bw), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 600 mg/kg bw), MEPOS (Methanolic extract of *Pandanus odoratissimus* seed 900 mg/kg bw).

DISCUSSION

management of diabetes mellitus with a substance with no side effect is still a challenge in the medical system. Consequently, there is an increase in the demand for natural products with anti-hyperglycemic activity but with fewer or no side effect. It is generally known that medicinal plants are used for treating diabetes due to their effectiveness, safety and acceptability. Most of these medicinal plants have antidiabetic in nature and contain different substances reported for various activities (30). Some reports showed that phytochemicals are responsible for some anti-hyperglycemic and hyperlipidemic activities (31). The phytochemical test showed the presence of flavonoid, tannins, saponin and quinons in the seed extract of *P. odoratissimus*. Also, it has been shown that flavonoid play a vital role in controlling glucose level and lipid concentration. These compounds are also known to regenerate damaged β cells in alloxan induce animal and act as insulin secretagogues, an important medicine for type 2 diabetes, as well as help in the secretion of insulin, thereby maintaining the blood glucose level (32). Other components such as tannins and saponin have been found to hinder glucose transport by inhibiting sodium glucose cotransporter 1 (S-GLUT - 1) in intestinal brush border cells (66). In addition, saponins are known to lower cholesterol in intestinal lumen, preventing its absorption and/or binding to bile acids, causing a reduction in the entero hepatic circulation of bile acids, and increasing fecal excretion (34). Increased bile acid excretion is offset by its synthesis from cholesterol in the liver, thereby lowering the plasma cholesterol (34, 35). It has also been reported that the administration of polyphenolic compound to alloxan-induced diabetic rats reduced hyperlipidemia and this was attributed to reduction in the activity of hepatic HMG-CoA reductase, which is the first committed enzymatic step in cholesterol synthesis (36). This lowers elevated LDL cholesterol levels, hence, reducing the common coronary events and death from coronary heart disease (CHD) which occurs in diabetic

patients (37). Therefore, the observed hypolipidemic effect of *P. odoratissimus* seed could be linked to the synergistic action of phytochemicals like saponin and polyphenolic compound in it. Additionally, fiber of plants interfere with carbohydrate absorption, thereby maintaining the blood glucose level.

Alloxan is a selective β cytotoxic agent widely used to mimic diabetic pathology rodent (38). It is uric acid derivative which causes significant reduction in serum insulin, thus, destroying the β cells of the islets of Langerhans (39) and causing hyperglycemia in rats (40). In addition, it is selectively toxic for pancreatic β cells of the islets of langerhans thereby inducing necrosis (41). Alloxan is cytotoxic glucose analog with a molecular shape similar with glucose (42) and has two pathological effects, namely, selective reduction of glucose-induced insulin secretion through specific inhibition of glucokinase and generation of free radicals. The mechanism through which alloxan induce diabetes is the generation of free radicals which damage and cause the death of β cells; it also infiltrate to the pancreatic β cells through the GLUT 2 transporter (43). Also, it is reduced in cytosol, in the presence of different cellular reducing agents to dialuric acid. This leads to a redox cycle and production of superoxide radicals (O_2^-), and the dismutation of these radicals generates hydrogen peroxide (H_2O_2) which reacts with ferrous (Fe^{2+}) to form hydroxyl radical ($OH\cdot$), a high oxidative agent (40). These oxidative agents cause necrosis in pancreas β cells, thereby leading to diabetes mellitus type 1 (38). Insulin is secreted by pancreas, which is the primary organ involved in determining the organism's dietary and energetic status through the amount of glucose present in the blood (44). This study shows that the *P. odoratissimus* seed extract reduced the glucose level in animal made diabetic with alloxan. One possible explanation for this could be the presence of antioxidant in the plant part, as reported in a previous study (45). Antioxidants are protective agents which inactivate the ROS causing cell damage (46). Interestingly, plants with flavonoids possess

both antioxidant and anti-hyperglycemic activities (47). Additionally, it has been reported that the antioxidant activity of herbal hypoglycemic plant extract could form a protective mechanism against the ROS associated with chronic hyperglycemia and diabetic complications such as microvascular and macrovascular conditions (46). Also, albino rats have surviving β cells, capable of regenerating due to the use of lower dose alloxan (125 mg/kg body weight), which produce partial destruction of pancreatic β cells (48). Other possible mechanism include the stimulation of cells and subsequent release of insulin and activation of the insulin receptor. Estimation of insulin level and receptor might give more insight into the mechanism of antidiabetic activity exhibited by the plant extract (49).

Another important clinical marker in diabetes is the glycosylate haemoglobin. It helps to regulate the degree of protein glycation during diabetes mellitus. In insistent hyperglycemic, formation of HbA1c takes place through the non-enzymatic reaction between free amino groups of haemoglobin and glucose. In diabetic patients, HbA1c levels help to assess both the long-term glycaemic control and the progression of damage or development of complication associated with the condition. He marked increase in HbA1c levels was noticed in the alloxan-induced diabetic animals, compared with the control group. However, treatment with the seed extract showed a reduction in HbA1c levels. This could be as a result of the reduction in the blood glucose caused through increasing insulin secretion by rejuvenated pancreatic β cells (50).

One of the complications as diabetic hyperglycemia is dyslipidemia. Generally, lipid play a major role in maintaining the integrity, structure, as well as the function of the biomembrane. However, due to insulin deficiency, alteration in lipid and lipoprotein associated with diabetes have a higher chance of developing into atherosclerosis. This dyslipid profile during diabetes, include increase in the level of total cholesterol (TC), triglyceride (TG), low density lipoprotein (LDL) and decrease in plasma high density lipoprotein (HDL) (51). The serum lipid profile of rats was evaluated in this study and as expected, untreated diabetic animals showed a significant increase in TG and LDL concentration but with low level of HDL. This increase in serum lipid is mainly due to the increase in acid mobilization from adipose tissue, since insulin inhibits 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase, which is the key enzyme in cholesterol biosynthesis (52). Hence, insulin deficiency or resistance could be responsible for hyperlipidemia. Also, treatment of type 2 diabetic rats with the methanolic seed extract (900 mg/kg body weight), completely reversed dyslipidemia as shown by the significant decrease ($p < 0.05$) in TC, TG and LDL, coupled with the increase in HDL ($p < 0.05$). These alleviating effects, therefore, show the antihyperlipidemic potential of methanolic extract of *P. odoratissimus* seeds.

CONCLUSION

The beneficial effects on the regulation of serum glucose levels and hyperlipidemia in alloxan-induced diabetic rats is evident in treatment with the dose of 900 mg/kg body weight of the *P. odoratissimus* seed extract. There is need for further research on it which could possibly be used in the development of diabetic drugs. Based on the results, the seed extract was

more potent and effective than the standard drug. Therefore, this study give scientific support to the use of *P. odoratissimus* in traditional beliefs for the treatment of diabetes and could be further explored for producing effective drugs for this condition due to its antidiabetic activity.

Acknowledgment

The authors are grateful to the Faculty of Biology, Universitas Nasional Jakarta, Indonesia for providing space in the animal house.

Conflict of interest

The authors reported no conflict of interest

REFERENCES

1. Cho NH. Key points for IDF Diabetes Atlas 2017. International Diabetes Federation Congress 2017, 4-8 December, Abu Dhabi. IDF Diabetes Atlas, 8th ed. 2017; p., 17-18.
2. American Diabetes Association (ADA). Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2013; 36: 67-74. PMID: 23264425 ; PMCID: PMC3537273.
3. Anilkumar MD, Naseeruddin MI. Protective effects of *Andrographis paniculata* against endothelial dysfunction in diabetic wistar rats. *J Pharmacol Toxicol* 2008; 3:311-17. doi : 10.3923/jpt.2008.311.317.
4. Schwartz SL. Diabetes and dyslipidemias. *Diabetes Obes Metab*. 2006; 8:355-64. PMID:16776742 doi: 10.1111/j.1463-1326.2005.00516.x
5. Balassa EO. Hyperlipidaemia in diabetes. *Medicographia* 1985; 7:11-14.
6. Davis SN : Insulin, oral hypoglycemic agent and the pharmacology of the endocrine pancreas. In *Textbook of The Pharmacological Basis of Therapeutics* Edited by Bruton LL, Lazo JS, Parker KL. New York: Mc Graw Hill; 5th Edition, 2006:1613-1645.
7. Inzucchi, SE 2002. Oral antihyperglycemic therapy for type 2 diabetes mellitus. *Scientific Review J Am Med Assoc* 287: 360-372. doi:10.1001/jama.287.3.373.
8. Halim EM, Hussain A. Hypoglycemic, hypolipidemic and antioxidant properties of combination of curcumin from *Curcuma longa*, Linn, and partially purified product from *Abroma augusta*, Linn. in streptozotocin induced diabetes. *Ind J Clin Biochem* 2002; 17:33-43. doi: 10.1007/BF02867969; PMCID: PMC3454113; PMID: 23105348.
9. Ghorbani A. Best herbs for managing diabetes : A review of clinical studies. *Braz J Pharm Sci* 2013; 49:413-422. doi: 10.1590/S1984-82502013000300003.
10. Oudhia P. *Pandanus odoratissimus* L.f. Record from PROTA4U. Brink, M. & Achigan-Dako, E.G. (Editors). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands. 2011. <http://www.prota4u.org/search.asp>. Accessed 1 March 2020.
11. Marpaung DWAK, Pasaribu N and Athorick TA. Taxonomy study Of *Pandanus* (Pandanaceae) in swamp area, Aceh Singkil. 2013; *Jurnal Natural* 13(2): 55-63.

12. Adkar PP, Jadhav PP, Ambavade SD, Sheike TT, Bhaskar VH. Protective effect of leaf extract of *Pandanus odoratissimus* Linn on experimental model of epilepsy. *Int J Nutr Pharmacol Neuro Dis* 2014; 4: 81-31. doi:10.4103/2231-0738.129590.
13. Kumar S, Dagar S, Kumar P, Kumar S, Kumar D. Antifertility effect of hydroalcoholic extract of *Pandanus odoratissimus* L. Leaves. *Porto Biomed* 2017; 5:167-169. <http://dx.doi.org/10.1016/j.pbj.2017.03.00>.
14. Ramesh L, Abhaykumar K, Reddy VC. 2010. Anti-inflammatory activity of *Pandanus odoratissimus* extract. *Int J Pharmacol* 2010; 6: 311-314. doi: 10.3923/IJP.2010.311.314.
15. Raju S, Subbaioah NV, Reddy KS, Das A, Murugan KB 2011. Potential of *Pandanus odoratissimus* as CNS depressant in Swiss albino mice. *Braz J Pharm Sci* 201; 47(3): 629-634.
16. Raj GG, Varghese HS, Kotagiri S, Swamy BMW, Swamy A, Pathan RK. Anticancer studies of aqueous extract of root and leaves of *Pandanus odoratissimus* f.feren. (Y.Kimura) Hatus: an in vitro approach. *J Trad Complement Med* 2014; 4(4): 279-284. doi: 10.4103/2225-4110.129199.
17. Venkantesh R, Kusuma R, Sateesh V, Madhara RB, Ramesh M, 2012. Antidiabetic activity of *Pandanus odoratissimus* root extract. *Ind J Pharm Edu Res* 2012; 46 (4): 340-345.
18. Suvetha SP, Kumar TS, Kumaraesan K, Raphael VS, Muthukumara V, Thirugnanan N. 2018. *In vitro* therapeutic correlative studies of *Pandanus odoratissimus* flowers against diabetic foot ulcer-causing bacterial pathogens. *Int J Green Pharm* 2018; 12(3): 601-606.
19. Nahid S, Mohajon M, Thousif A Md, Paul A, Barua A and Islam Md S. Phytochemical and Pharmacological screening of methanolic extract and its subfraction of *Pandanus odoratissimus* Linn.Fruits. *Int J Res Pharm Biosci* 2019; 6(12): 5-15.
20. Tiwari, AK, Rao JM. Diabetes Mellitus and Multiple Therapeutic Approaches of Phytochemicals: Present Status and Future Prospects. *Curr Sci* 2002; 83: 30-38. www.jstor.org/stable/24106071.
21. Hakkim FL, Girija S, Kumar RS, Jalaluddeen MD. Effect of aqueous and ethanol extracts of *Cassia auriculata* L. Flowers on diabetes using alloxan induced diabetic rats. *Int J Diabetes Metab* 2007; 15: 100-106.
22. Ng MY, Peng CH, Chan KC, Yang YS, Huang CN, Ng CJ. The hypolipidemic effect of *Hibiscus tariffa* polyphenols via inhibiting lipogenesis and promoting hepatic lipid clearance. *J Agric Food Chem* 2010;58: 850-859. doi: 10.1021/jf903209w. PMID:20017484.
23. Osibemhe M, Onoagbe IO. Qualitative and quantitative phytochemical evaluation of *Strophatus hispidus* stem bark. *IOSR J Pharm Biol Sci* 2015;10(2): 124-128. doi: 10.9790/3008-1021120124.
24. Akinyemi KO. Screening of crude extracts of six medicinal plants used in south-west Nigerian unorthodox medicine for anti-methicilin resistant *Staphylococcus aureus* activity. *BMC Complement Altern Med* 2005: 2205: 1-7. doi:10.1186/1472-6882-5-6.
25. Bendiabdellah A, Dib MA, Meliani N, Djabou N, Allah H, Tabti B. Preliminary phytochemical screening and antioxidant activities of solvent extract from *Daucus crinitus* Desf., from Algeria. *J Appl Pharm Sci* 2012; 2(7): 92-95. doi: 10.7324/JAPS.2012.2710.
26. De S, Dey YN, Ghosh AK 2010. Phytochemical investigation and chromatographic evaluation of different extracts of tuber of *Amorphophallus paenifolius* (Araceae). *Int J Pharmaceu Biomed Res* 2010; 1: 150-157.
27. Braca A, Tommasi ND, Bari LD, Pizza C, Politi M, Morelli I. Antioxidant principles from *Bauhinia terapotensis*. *J Nat Prod* 2001; 64: 892-895. doi: 10.1021/np0100845 PMID: 11473417.
28. Viturro C, Molina A, Schmeda-Hirschmann G. Free radical scavengers from *Mutisia friesiana* (Asteraceae) and *Sanicula graveolens* (Apiaceae). *Phytother Res* 1999; 13:422-424. DOI: 10.1002/(sici)1099-1573(199908/09)13:5<422::aid-ptr462>3.0.co;2-m ; PMID: 10441784.
29. Oleszek M, Kozachok S. Antioxidant activity of plant extract and their effect on methane fermentation in bioreactors. *Int Agrophys* 2018; 32: 395-401. doi: 10.1515/intag-2017-0031.
30. Loew D, Kaszkin M. Approaching the problem of bioequivalence of herbal medicinal products. *Phytother Res* 2002; 16:705-711. doi: 10.1002/ptr.1248 ; PMID: 12458470.
31. Kumar R, Pate DK, Prasad SK, Sairam K, Hemalatha S. Antidiabetic activity of alcoholic leaves extract of *Alangium lamarkii* Thwaites on streptozotocin-nicotinamide induced type 2 diabetic rats. *Asian Pac J Trop Med* 2011; 4: 904-909. doi: 10.1016/S1995-7645(11)60216-2.
32. Jung M, Park M, Lee HC, Kang YH, Kang ES, Kim SK. Antidiabetic agents from medicinal plants. *Curr Med Chem* 2006;13(10):1203-18. doi: 10.2174/092986706776360860 ; PMID: 16719780.
33. Petchi RR, Parasuraman S, Vijaya C. Antidiabetic and antihyperlipidemic effects of an ethanolic extract of the whole plant of *Tridax procumbens* (Linn.) in streptozotocin-induced diabetic rats. *J Basic Clin Pharm* 2013; 4: 88-92. doi:10.4103/0976-0105.121655; PMID:24808679.
34. Timi SO, Omotosho OE, Roimi OA. Persistence of insulin resistance in alloxan induced diabetic rats treated with the juice of *Asystasia gangetica* leaves. *Phcog Mag* 2011; 7: 25-30. doi: 10.4103/0973-1296.75887 ; PMID:21472075.
35. James DB, Owolabi OA, Ibrahim AB, Folorunsho DF, Bwalla I, Akanta F. 2010. Changes in lipid profile of aqueous and ethanolic extract of *Bleighias apida* in rats. *Asian J Med Sci* 2010; 2:177-180.

36. Kumarappen CT, Rao TN, Mandal SC. Polyphenolic extract of *Ichnocarpus frutescens* modifies hyperlipidemia status in diabetic rats. *J Cell Mol Bio* 2007; 6: 175-185.
37. Richard AH, Pamela CC. Lippincott's Illustrated Reviews: Pharmacology Lippincott Williams and Wilkin. 4th Edn., A Wolters Kluwer Company, Baltimore, 2009; pp: 249-295.
38. Lenzen S. The mechanisms of alloxan- and streptozotocin induced diabetes. *Diabetologia* 2008; 51(2): 216-226. doi: 10.1007/s00125-007-0886-7 ; PMID: 18087688.
39. Patel DK, Kumar R, Laloo D, Hemalatha S. Diabetes mellitus: an overview on its pharmacological aspects and reported medicinal plants having antidiabetic activity. *Asian Pac J Trop Bio* 2012; 2 (5): 411-420. DOI: 10.1016/S2221-1691(12)60067-7 ; PMID: PMC3609313 PMID: 23569941.
40. Szkudelski T. 2001. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol Res* 2001; 50: 537-546. PMID: 11829314.
41. Etuk EU. 2010. Animals models for studying diabetes mellitus. *Agric Biol J N Am* 2010; 2:130-134.
42. Berraouan A, Ziyat A, Moudi H, Legssyer A, Mohammed A, Bnouham M. Evaluation of protective effect of cactus pear seed oil (*Opuntia ficus-indica* L. Mill.) against alloxan induced diabetes in mice. *Asian J Trop Med* 2015; 8: 532-537. doi: 10.1016/j.apjtm.2015.06.013; PMID: 26276283.
43. Elsner M, Tiedge M, Guldbakke B, Munday R, Lenzen S. Importance of the GLUT2 glucose transporter for pancreatic beta cell toxicity of alloxan. *Diabetologia* 2002; 45:1542-1549. doi: 10.1007/s00125-002-0955-x; PMID: 12436338.
44. Edem DO. Hypoglycemic effects of ethanolic extracts of alligator pear seed (*Persea americana* Mill) in rats. *Eur J Sci Res* 2009; 33(4): 669-678.
45. Queiroz FE, Kuhl C, Terreaux C, Mavi S, Hostettmann K. New dihydroalkylphenones from *Lansea edulis*. *Nat Prod* 2003; 66: 578-580. doi: 10.1021/np0202428 ; PMID: 12713423.
46. Joseph SA, Theophilus FC, Dominic EA, Rita DA, Kwaw ML, Kofi A, et al. Antihyperglycaemic and antioxidant effects of *Adenia lobata* engl. (Passifloraceae) in streptozotocin-induced diabetic rats. *Afr J Tradit Complement Altern Med* 2013; 10: 386-393. doi: 10.4314/ajtcam.v10i5.28; PMID: PMC3847435 PMID: 24311855.
47. Petchi RR, Parasuraman S, Vijaya C. Antidiabetic and hyperlipidemic effects of an ethanolic extract of the whole plant of *Tridax procumbens* (Linn.) in streptozotocin-induced diabetic rats. *J Basic Clin Pharm* 2013; 4:88-92. doi: 10.4103/0976-0105.121655 ; PMID: PMC3979266 ; PMID: 24808679.
48. Far MJ, Sanchez R, Grau AN, Sanchez SS. Hypoglycemic effect of the water extract of *Smalanthus sonchifolius* (yacon) leaves in normal and diabetic rats. *J Ethnopharmacol* 2001; 74:125-132. doi: 10.1016/S0378-8741(00)00351-2 ; PMID: 11167030.
49. Mandal SC, Maity TK, Das J, Pal M, Saha BP. Hepatoprotective activity of *Ficus racemosa* leaf extract on liver damage caused by carbon tetrachloride in rat. *Phytother Res* 1999; 13(5):430-32. DOI: 10.1002/(sici)1099-1573(199908/09)13:5<430::aid-ptor56>3.0.co;2-g ; PMID: 10441787.
50. Aboonabi A, Rahmat A, Othman F. Anti-oxidant effect of pomegranate against streptozotocin-nicotinamide generated oxidative stress induced diabetic rats. *Toxicol Rep* 2014; 1: 915-22. doi: 10.1016/j.toxrep.2014.10.022 ; PMID: PMC5598111 ; PMID: 8962304.
51. Grundy SM, Benjamin IJ, Burke GL, Chait A, Eckel RH, Howard BV et al. Diabetes and cardiovascular disease: a statement for healthcare professionals from the American Heart Association. *Circulation* 1999; 100:1134-46. doi: 10.1161/01.cir.100.10.1134 ; PMID: 10477542.
52. Betteridge J. Lipid disorders in diabetes mellitus. In *Textbook of Diabetes*. Edited by Pickup J, Williams G. London: Blackwell Science; 2002:551-553.

Antihyperglycemic and Antihyperlipidemic Effects of Methanolic Seeds Extract of Pandanus odoratissimus in Alloxan-Induced Diabetic Rats

ORIGINALITY REPORT

44%

SIMILARITY INDEX

37%

INTERNET SOURCES

38%

PUBLICATIONS

14%

STUDENT PAPERS

PRIMARY SOURCES

1	www.sysrevpharm.org Internet Source	4%
2	Loubna Ait Dra, Souad Sellami, Hanane Rais, Faissal Aziz, Abdallah Aghraz, Khalid Bekkouche, Mohamed Markouk, Mustapha Larhsini. "Antidiabetic potential of Caralluma europaea against alloxan-induced diabetes in mice", Saudi Journal of Biological Sciences, 2018 Publication	2%
3	bmccomplementmedtherapies.biomedcentral.com Internet Source	2%
4	www.frontiersin.org Internet Source	2%
5	www.iosrjournals.org Internet Source	1%
6	ukjpb.com Internet Source	1%

7	Submitted to Universitas Siliwangi Student Paper	1%
8	file.scirp.org Internet Source	1%
9	maxwellsci.com Internet Source	1%
10	www.scholarsresearchlibrary.com Internet Source	1%
11	banglajol.info Internet Source	1%
12	MR Saha, SMR Hasan, R Akter, MM Hossain, MS Alam, MA Alam, MEH Mazumder. " <i>In vitro</i> free radical scavenging activity of methanol extract of the leaves of <i>Mimusops elengi</i> Linn.", Bangladesh Journal of Veterinary Medicine, 1970 Publication	1%
13	www.ijddr.in Internet Source	1%
14	www.ncbi.nlm.nih.gov Internet Source	1%
15	Shahid Akbar. "Handbook of 200 Medicinal Plants", Springer Science and Business Media LLC, 2020 Publication	1%

16	www.thieme-connect.com Internet Source	1%
17	www.mdpi.com Internet Source	1%
18	www.tandfonline.com Internet Source	1%
19	Kousik Ahmed Khan, Manik Zahan, Fatema Zohura Talukder, Riaz Uddin et al. "Alternanthera bicolor Produces Hypoglycemic Effect in Alloxan-Induced Diabetic Mice through its Antioxidant Activity", Dhaka University Journal of Pharmaceutical Sciences, 2019 Publication	1%
20	ismj.bpums.ac.ir Internet Source	1%
21	www.informaticsjournals.com Internet Source	1%
22	Submitted to Tikrit University Student Paper	1%
23	www.arjournals.org Internet Source	1%
24	Shanmugam, K.R.. "Neuroprotective effect of ginger on anti-oxidant enzymes in streptozotocin-induced diabetic rats", Food and Chemical Toxicology, 201104	1%

25 worldwidescience.org 1%

Internet Source

26 Ramesh Londonkar, Abhaykumar Kamble, V. Chinnappa Reddy. "Anti-Inflammatory Activity of Pandanus odoratissimus Extract", International Journal of Pharmacology, 2010 <1%

Publication

27 link.springer.com <1%

Internet Source

28 www.ajol.info <1%

Internet Source

29 s3.amazonaws.com <1%

Internet Source

30 www.dovepress.com <1%

Internet Source

31 Arati Malpani, Nitin Mahurkar, Urmila Aswar. "Phytochemical analysis and antifertility potential of Cynodon dactylon in female Wistar rats: A herbal approach towards contraception", Chinese Herbal Medicines, 2020 <1%

Publication

32 Ivanildo I. da S. Júnior, Humberto de Moura Barbosa, Débora C. R. Carvalho, Ruideglan de Alencar Barros et al. "Brazilian Attenuated <1%

Hyperglycemia, Dyslipidemia, and Prooxidant Status in Alloxan-Induced Diabetic Rats ", The Scientific World Journal, 2017

Publication

33

www.usa-journals.com

Internet Source

<1%

34

japsonline.com

Internet Source

<1%

35

medsci.org

Internet Source

<1%

36

www.scielo.br

Internet Source

<1%

37

ijpsr.com

Internet Source

<1%

38

Ji Su Kim, Jung Bong Ju, Chang Won Choi, Sei Chang Kim. "Hypoglycemic and Antihyperlipidemic Effect of Four Korean Medicinal Plants in Alloxan Induced Diabetic Rats", American Journal of Biochemistry and Biotechnology, 2006

Publication

<1%

39

0-ftp.crossref.org.wam.leeds.ac.uk

Internet Source

<1%

40

www.degruyter.com

Internet Source

<1%

- 41 Imen Dahech, Karima Srih Belghith, Khaled Hamden, Abdelfattah Feki, Hafedh Belghith, Hafedh Mejdoub. "Antidiabetic activity of levan polysaccharide in alloxan-induced diabetic rats", International Journal of Biological Macromolecules, 2011
Publication <1%
-
- 42 www.ijsidonline.info
Internet Source <1%
-
- 43 L. Wu, H. Qiao, Y. Li, L. Li. "Protective roles of puerarin and Danshensu on acute ischemic myocardial injury in rats", Phytomedicine, 2007
Publication <1%
-
- 44 www.cmj.org
Internet Source <1%
-
- 45 Singh Nayak. "Evaluation of antidiabetic and antioxidant activity of aerial parts of Hyptis suaveolens Poit.", African Journal of Pharmacy and Pharmacology, 2013
Publication <1%
-
- 46 Edible Medicinal and Non Medicinal Plants, 2014.
Publication <1%
-
- 47 Rathee Permender, Chaudhary Hema, Rathee Sushila, Rathee Dharmender, Kumar Vikash. "Antidiabetic Potential of Fabaceae Family: An <1%

Overview", Current Nutrition & Food Science, 2010

Publication

48	www.japsonline.com Internet Source	<1%
49	article.jplantsciences.org Internet Source	<1%
50	raw.githubusercontent.com Internet Source	<1%
51	El-Batran, S.A.. "Studies on the glycemic and lipidemic effect of monopril and losartan in normal and diabetic rats", Pharmacological Research, 200408 Publication	<1%
52	Submitted to Universitas Nasional Student Paper	<1%
53	Submitted to University of KwaZulu-Natal Student Paper	<1%
54	www.mims.com Internet Source	<1%
55	researchspace.ukzn.ac.za Internet Source	<1%
56	Mohini C. Upadhye, Uday Deokate, Rohini Pujari, Vishnu Thakare. " Antidiabetic Effects of Ethanolic Extract of (L.) Roots ", Current	<1%

57

iraqijms.com

Internet Source

<1%

58

Bakhashwain , Amal Salem Ahmad. "The Antidiabetic Effect of Nigella Sativa and Propolis on Streptozotocin-Induced Diabetes and

Diabetic Nephropathy in Male Rats = تأثير مضاد السكري لحبة البركة والعكبر على ذكور الجرذان المصابة بالسكري", "المستحث باستريبتوزوتوسين ومعالجة اعتلال الكلى السكري",

King Abdulaziz University : Scientific Publishing Centre, 2016

Publication

<1%

59

elsevier.pt

Internet Source

<1%

60

K.L. Joy, R. Kuttan. "Anti-diabetic activity of Picrorrhiza kurroa extract", Journal of Ethnopharmacology, 1999

Publication

<1%

61

Bakhdar , Fatmah Abdullah Omar. "Effect of Histone Deacetylase Inhibitor (Vorinostat) on New-Onset Diabetes Induced by Treatment with

Tacrolimus = تأثير مثبطات الهيستون دياسيتايلاز

(فورينوستات) على مرض السكري الجديد الناتج عن العلاج

"بالتاكروليمس", King Abdulaziz University : Scientific Publishing Centre, 2020

Publication

<1%

62

creamjournal.org

Internet Source

<1%

63

www.hindawi.com

Internet Source

<1%

64

Paola Isabel Angulo-Bejarano, María del Rocío Gómez-García, Maria Elena Valverde, Octavio Paredes-López. "Nopal (*Opuntia* spp.) and its Effects on Metabolic Syndrome: New Insights for the Use of a Millenary Plant", *Current Pharmaceutical Design*, 2019

Publication

<1%

65

Bartholomew O. Ibeh, Maxwell I. Ezeaja. "Preliminary study of antidiabetic activity of the methanolic leaf extract of *Axonopus compressus* (P. Beauv) in alloxan-induced diabetic rats", *Journal of Ethnopharmacology*, 2011

Publication

<1%

66

eprints.covenantuniversity.edu.ng

Internet Source

<1%

67

academicjournals.org

Internet Source

<1%

68

www.science-truth.com

Internet Source

<1%

69

Samantha Beck, Heather Mathison, Toma Todorov, Esli-Armando Calderon-Juarez, Olga

<1%

R. Kopp. "A Review of Medicinal Uses and Pharmacological Activities of *Tridax Procumbens* (L.)", *Journal of Plant Studies*, 2018

Publication

70

ajp.mums.ac.ir

Internet Source

<1%

71

www.medsci.org

Internet Source

<1%

72

Folorunsho A. Ajayi, Olubukola. S. Olorunnisola, Adewale. Adetutu, Folashade G. Olorunfemi et al. "Anti-hyperglycaemic and Mode of Action of *Thaumatococcus danielli* (BENN.) BENTH Ethanol Leave Extract in Streptozotocin-induced Diabetic Rats", *Asian Journal of Research in Medical and Pharmaceutical Sciences*, 2019

Publication

<1%

73

www.kau.edu.sa

Internet Source

<1%

74

Edible Medicinal and Non Medicinal Plants, 2015.

Publication

<1%

75

journals.tbzmed.ac.ir

Internet Source

<1%

76

Tchamadeu, M.C.. "Hypoglycaemic effects of *Mammea africana* (Guttiferae) in diabetic rats",

<1%

- 77 Ufele Angela Nwogor. "Effects of Prolonged Administration of Aqueous Extract of *Ocimum gratissimum* (Scent Leaf) on Blood Glucose and Lipid Profile in Alloxan Induced Diabetic Albino Rats", American Journal of Biomedical and Life Sciences, 2016 <1%
- Publication
-

- 78 Abo Gazia, Maha M.. "Preclinical immunohistochemical biomarkers for testicular changes in hyperlipidemic rats and the role of *Ocimum gratissimum* :", The Egyptian Journal of Histology, 2013. <1%
- Publication
-

- 79 Leila Mousavi, Rabeta Mohd Salleh, Vikneswaran Murugaiyah, Mohd Zaini Asmawi. "Hypoglycemic and anti-hyperglycemic study of *Ocimum tenuiflorum* L. leaves extract in normal and streptozotocin-induced diabetic rats", Asian Pacific Journal of Tropical Biomedicine, 2016 <1%
- Publication
-

- 80 Gilani, A.H.. "Ethnopharmacological studies on antispasmodic and antiplatelet activities of *Ficus carica*", Journal of Ethnopharmacology, 20080902 <1%
- Publication
-

81

"Bioactive Natural Products for Pharmaceutical Applications", Springer Science and Business Media LLC, 2021

Publication

<1%

82

Xing, Ronge, Xiaofei He, Song Liu, Huahua Yu, Yukun Qin, Xiaolin Chen, Kecheng Li, Rongfeng Li, and Pengcheng Li. "Antidiabetic Activity of Differently Regioselective Chitosan Sulfates in Alloxan-Induced Diabetic Rats", Marine Drugs, 2015.

Publication

<1%

83

Rahmat Khan, Wasim Ahmad, Mushtaq Ahmed. "Stellaria media attenuates the hyperglycemia and hyperlipidemia in alloxan-induced diabetic rat", Bangladesh Journal of Pharmacology, 2019

Publication

<1%

84

Abdullah Zehad, Gazi Jahirul Islam, Mamunur Rashid, Nusrat Jahan Juthy, Shaheda Zannah. "Antidiabetic and Antihyperlipidemic Activities of Methanolic Leaf Extract of *Stephania japonica* in Alloxan Induced Diabetic Rats", Pharmacology & Pharmacy, 2017

Publication

<1%

85

Deepti Dabas, Rachel Shegog, Gregory Ziegler, Joshua Lambert. "Avocado (*Persea americana*) Seed as a Source of Bioactive Phytochemicals",

<1%

Current Pharmaceutical Design, 2013

Publication

86

Edible Medicinal And Non-Medicinal Plants,
2014.

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On